

I.L. Stadnik<sup>1</sup>, F.Zh. Abilkanova<sup>1</sup>, Ye.V. Kudryavtseva<sup>1</sup>, S.N. Nikolskiy<sup>1,2</sup>, A.S. Masalimov<sup>1</sup>

<sup>1</sup>Karagandy University of the name of academician E.A. Buketov, Kazakhstan;

<sup>2</sup>Altay State University, Barnaul, Russia

(\*Corresponding author's e-mail: i.stadnik@mail.ru)

## ESR-study of the proton exchange with aliphatic amino acids in toluene


This paper presents the results of an ESR spectroscopic study of the intermolecular proton exchange kinetics with some isomers of aminocaproic acid, such as 2-aminohexanoic and 6-aminohexanoic acids in the toluene indifferent medium. A stable semiquinone radical, namely 3,6-di-*tert*-butyl-2-hydroxyphenoxy, was used as a spin probe. The ESR spectra were recorded on a RE-1306 spectrometer. The obtained dynamic spectra of the intermolecular proton exchange process are given in this article. Modeling of the proton exchange ESR spectra was carried out using a previously developed program, which is based on a four-jump model of this reaction and modified Bloch equations. The kinetic parameters of the process of acid-base interaction of the spin probe with amino acids and the activation barrier of the reaction were estimated with a minimum error based on a comparison of the model spectra with the experimental ones. The obtained data were analyzed and compared with previously known information. It was found that the reaction rate of the intermolecular proton exchange between 3,6-di-*tert*-butyl-2-hydroxyphenoxy and studied aliphatic amino acids is comparable to the same with aliphatic carboxylic acids. But in our case, there is an increase in the activation barrier, which apparently is associated with a negative effect on the intramolecular hydrogen bonding process between the amino and the carboxyl groups in the amino acid molecule.

**Keywords:** ESR-spectroscopy, proton exchange reactions, spin probe, semiquinone radical, Bloch equation, aminohexanoic acid, 3,6-di-*tert*-butyl-2-hydroxyphenoxy, OH-acids.


### Introduction

The acid-base interactions are very common phenomena in nature. They are widely used in scientific and practical activities. Theoretical ideas about acids and bases are often of primary importance in the formation of all conceptual systems of chemistry, and have a versatile influence on the development of many theoretical concepts in all fundamental chemical disciplines. Typical acid-base interactions according to the Bronsted-Lowry theory include protolytic reactions involving proton. The proton exchange reaction is one of the varieties of such reactions. To date, a large number of papers have been published on the above acid-base interactions. However, as can be seen from the literature analysis, the most studies of protolytic reactions have been carried out in aqueous media, and there are few data on such studies in non-aqueous media [1–6].

Previously, we have carried out ESR spectroscopic studies of proton exchange reactions between a stable semiquinone radical as a spin probe and various organic acids in non-aqueous media [7, 8]. It seems interesting to study the proton exchange reaction with aminohexanoic acids (1), namely 2-aminohexanoic ( $\alpha$ -aminocaproic) and 6-aminohexanoic ( $\epsilon$ -aminocaproic):



2-aminohexanoic acid



6-aminohexanoic acid

(1)

This study will make it possible to determine the protolytic ability of these acids and to evaluate the influence of their structure on the rate of the proton exchange reaction.

Amino acids play an important role in various biochemical processes. The interest of scientists in this class of compounds is very great. This is confirmed by a large number of publications devoted to the study of their physicochemical properties [9–14].

### Experimental

The method of recrystallization from tetrahydrofuran was used to purify 3,6-di-*tert*-butylcyclohexa-3,5-diene-1,2-dione (3,6-di-*tert*-butyl-*o*-benzoquinone) (N.D. Zelinsky Institute of Organic Chemistry of the Russian Academy of Sciences, >98 % purity) and 3,6-di-*tert*-butylbenzene-1,2-diol (3,6-di-*tert*-butylcatechol) (N.D. Zelinsky Institute of Organic Chemistry of the Russian Academy of Sciences, >98 % purity).

Toluene (CJSC "Ekos-1", >98 % purity) was purified by distillation under atmospheric pressure over sodium hydroxide.


Amino acids, namely 2-aminohexanoic and 6-aminohexanoic (AppliChem GmbH), were used without additional purification, since their purity exceeded 99 %. Amino acids were placed in the desiccator for 48 hours before use [15].

The studied system of 3,6-di-*tert*-butyl-2-hydroxyphenoxy — amino acids was obtained by direct mixing of 3,6-di-*tert*-butylcyclohexa-3,5-diene-1,2-dione with 3,6-di-*tert*-butylbenzene-1,2-diol in a glass ampoule followed by the addition of an amino acid solution in toluene of a given concentration. Then the ampoule was degassed and sealed. The spectra were recorded on a RE-1306 spectrometer (Russia) with a unit for temperature control in the range of 200-400K. The spectrometer was upgraded with a device that converts the analog signal to digital. The ESREXCHANGE program was used to process the experimental data. Previously, we have developed this program in the FORTRAN algorithmic language [16]. The ESREXCHANGE program includes calculation of the theoretical spectrum, comparison of this spectrum with the experimental spectrum, calculation of the kinetic parameters of the intermolecular proton exchange reaction, Least-squares processing with determination of the standard error of measurement.

### Results and Discussion

The ESR spectrum of 3,6-di-*tert*-butyl-2-hydroxyphenoxy is a triplet of doublets. It is formed by the interaction of an unpaired electron with ring protons ( $a_H = 0.392$  mT) and with a proton of the hydroxyl group ( $a_{OH} = 0.162$  mT). The advantage of the spin probe used is the hyperfine structure of the radical's ESR spectrum, which is very sensitive to acid-base interactions [7, 8].

The process of intermolecular proton exchange with H-acids can be described by a four-jump model:



Along with the above reaction ( $A \leftrightarrow C$ ,  $B \leftrightarrow D$ ), it includes tautomeric transformations of a semiquinone radical, i.e. intramolecular migration of the hydrogen atom of the hydroxyl group between oxygen atoms ( $A \leftrightarrow B$ ,  $C \leftrightarrow D$ ) [17].

This four-jump model (2) and the modified Bloch equation formed the basis of the previously developed program for modelling intermolecular proton exchange. In this program, the ESR spectra are calculated from the coupling constants, rate constants, and linewidths of all chemical configurations of 3,6-di-*tert*-butyl-2-hydroxyphenoxy. The rate constants of the reaction of intermolecular proton exchange, the activation energy of the process, and the measurement error can be found by comparing the model spectra with the

experimental ones. It should be noted that the proton exchange process presumably has a “cooperative” mechanism, that is, a cyclic hydrogen bonded complex is formed during the reaction. The lifetime of this complex directly affects the rate of the whole process [17].

The ESR spectra of the 3,6-di-*tert*-butyl-2-hydroxyphenoxy — aminohexanoic acid system were obtained by the ESR method. The broadening of the hyperfine structure lines was noted with increasing temperature. This is characteristic of proton exchange and confirms that an intermolecular reaction of proton exchange is observed in the system under study.

For example, Figure 1 shows experimental and simulated ESR spectra of the 3,6-di-*tert*-butyl-2-hydroxyphenoxy — 6-aminohexanoic acid system in toluene.




Figure 1. ESR spectra of 3,6-di-*tert*-butyl-2-hydroxyphenoxy — 6-aminohexanoic acid (0.05 mol/l) system in toluene at different temperatures. Blue line is an experimental spectrum; red line is a simulated spectrum

The decrease in the central components of the ESR spectrum triplet (Fig. 1) indicates a slowdown in proton migration between oxygen atoms in the radical. This means that the resulting intermediate of semiquinone radical with acid is sufficiently stable.

The Arrhenius plot showed a good linear relationship between  $\lg k_{\text{exch}}$  and  $1/T$  (Fig. 2). The plot in Figure 2 also shows the values of the standard measurement error calculated by ESREXCHANGE program.




Figure 2. Arrhenius plot of the intermolecular rate constant ( $k_{\text{exch}}$ ) for 3,6-di-*tert*-butyl-2-hydroxyphenoxyl — 6-aminohexanoic acid system in toluene obtained from the experimental and theoretical ESR spectra

Kinetic parameters of intermolecular proton exchange reaction between 3,6-di-*tert*-butyl-2-hydroxyphenoxyl and aminohexanoic acids were calculated using the ESREXCHANGE program. The obtained data and measurement error are presented in Table 1.

Table 1

**Kinetic parameters of the intermolecular proton exchange reactions between 3,6-di-*tert*-butyl-2-hydroxyphenoxyl and H-acids in toluene solution**

H-acid	$k_{\text{exch}}$ (293K), l/mol's	$k_{\text{exch}}^0$ , l/mol's	$E_a$ , kJ/mol	pK <sub>a</sub>
2-Aminohexanoic acid	$(3.58 \pm 0.44) \cdot 10^7$	$1.96 \cdot 10^{10}$	$15.4 \pm 0.7$	2.34 9.83
6-Aminohexanoic acid	$(2.90 \pm 0.23) \cdot 10^7$	$5.95 \cdot 10^{10}$	$18.6 \pm 3.6$	4.43
Hexanoic acid	$(8.01 \pm 0.24) \cdot 10^7$	$2.78 \cdot 10^9$	$8.7 \pm 0.5$	4.88
Hexanedioic acid	$(7.22 \pm 0.70) \cdot 10^8$	$5.09 \cdot 10^9$	$4.8 \pm 0.8$	4.43 5.28
2-Anilinobenzoic acid	$(3.15 \pm 0.20) \cdot 10^7$	$9.25 \cdot 10^9$	$13.8 \pm 1.1$	5.28
Dodecan-1-amine	$(1.02 \pm 0.02) \cdot 10^9$	$3.64 \cdot 10^9$	$3.1 \pm 0.2$	10.63

Table 2 shows similar parameters with hexanoic, hexanedioic, 2-anilinobenzoic acids and dodecan-1-amine for comparison [17, 18]. Analysis of available data suggests that despite their amphoteric nature 2- and 6-aminohexanoic acids in toluene react with the spin probe at the rate comparable to the rate of the proton exchange reaction with aliphatic monocarboxylic acids (hexanoic acid is given for an example). The value of this rate is far from the rates with aliphatic amines [18]. Therefore, a proton of the carboxyl group of aminohexanoic acid is involved in the proton exchange. It is assumed that the proton exchange will occur both in the amino group and in the carboxyl group simultaneously. At the same time, an increase in the rate should be recorded at least by an order of magnitude, as in the case of dicarboxylic hexanedioic acid, but we do not observe this. It should be noted that the high activation energy of the process is a feature of the proton exchange reaction with 2- and 6-aminohexanoic acids. Apparently, there are competing reactions of the intra- and intermolecular interaction of the amino group with the carboxyl group of aminohexanoic acid. Also, this interaction is more pronounced in 6-aminohexanoic acid, where these functional groups occupy terminal positions, and the flexibility of the carbon skeleton of the molecule makes it possible to form fairly stable structures. All these processes have a negative effect on the resulting intermolecular complex with the spin probe and lead to an increase in the activation barrier. A similar situation is observed in the reaction of 3,6-di-*tert*-butyl-2-hydroxyphenoxyl with 2-anilinobenzoic acid.

### Conclusions

The intermolecular proton exchange reaction of the spin probe, namely 3,6-di-*tert*-butyl-2-hydroxyphenoxy, with 2-aminohexanoic and 6-aminohexanoic acids was carried out with the carboxyl group participation. The protolytic ability of aminohexanoic acid isomers in the toluene is comparable to that of aliphatic monocarboxylic acids and aromatic amino acids. The high activation barrier of the intermolecular proton exchange is due to the presence in the system of intramolecular and intermolecular interactions between the carboxyl and amino groups, which are typical for amino acids. This has a destructive effect on the intermediate reaction complex.

### References

- 1 Wang C.H. A comparative study on intermolecular hydrogen bond interactions in molecular dimers using different levels of computational methods / C.H. Wang, R.Q. Zhang, Z.J. Lin // *Journal of Theoretical & Computational Chemistry*. — 2012. — № 11(6). — P. 1237–1259. <https://doi.org/10.1142/s0219633612500836>
- 2 Bureiko S.F. The kinetic spectroscopy of hydrogen exchange processes in systems with hydrogen bond / S.F. Bureiko // 16th International Conference on Spectroscopy of Molecules and Crystals; Minist E., Sci Ukraine. — T. 5507: Proceedings of the Society of Photo-Optical Instrumentation Engineers (Spie) – Sevastopol, UKRAINE, 2003. — P. 143–153.
- 3 Limbach H.H. Double Proton Tautomerism via Intra- or Intermolecular Pathways? The Case of Tetramethyl Reductic Acid Studied by Dynamic NMR: Hydrogen Bond Association, Solvent and Kinetic H/D Isotope Effects / H.H. Limbach, S. Baumgartner, R. Franke, F. Mannle, G. Scherer, G.S. Denisov // *Molecules*. — 2021. — T. 26, № 14. <https://doi.org/10.3390/molecules26144373>
- 4 Grabowski S.J. Reaction pathway of proton transfer from the neutral to zwitterionic forms of amino acids. Support for a water molecule-mediated mechanism / S.J. Grabowski, T.M. Krygowski, B. Stepien // *Journal of Physical Organic Chemistry*. — 2000. — № 13(11). — P. 740–744. [https://doi.org/10.1002/1099-1395\(200011\)13:11<740::aid-poc304>3.3.co;2-t](https://doi.org/10.1002/1099-1395(200011)13:11<740::aid-poc304>3.3.co;2-t)
- 5 Wu Z.C. Proton affinity of arginine measured by the kinetic approach / Z.C. Wu, C. Fenselau // *Rapid Communications in Mass Spectrometry*. — 1992. — № 6(6). — P. 403–405. <https://doi.org/10.1002/rcm.1290060610>
- 6 Bureiko S.F. Structure and intermolecular interactions in complexes with hydrogen bond of a series of bifunctional nitrogen-containing compounds / S.F., Bureiko, S.Y. Kucherov // *Journal of Structural Chemistry*. — 2009. — № 50(4). — P. 712–721. <https://doi.org/10.1007/s10947-009-0109-9>
- 7 Masalimov A.S. EPR-spectroscopy of the fast proton exchange reactions in solutions / A.S. Masalimov, A.F. Kurmanova, A.U. Ospanov, A.A. Tur, S.N. Nikolskiy // *Bulletin of the Karaganda university — Chemistry*. — 2014. — № 73. — P. 30–35.
- 8 Nikolskiy S.N. Investigation of intermolecular proton exchange 3,6-di-*tert*-butyl-2-oxyphenoxy with phenol by ESR spectroscopy method / S.N. Nikolskiy, A.A. Tur, A.A. Yelchibekova, K.Zh. Kutzhanova // *Bulletin of the Karaganda university — Chemistry*. — 2015. — № 77. — P. 47–50.
- 9 Rozman M. Proton affinity of several basic non-standard amino acids / M. Rozman // *Chemical Physics Letters*. — 2012. — № 543. — P. 50–54. <https://doi.org/10.1016/j.cplett.2012.06.048>
- 10 Zhao G.D. Amino acid clusters supported by cellulose nanofibers for proton exchange membranes / G.D. Zhao, X.L. Xu, Y.B. Di, H. Wang, B.W. Cheng, L. Shi., Y. Yin // *Journal of Power Sources*. — 2019. — № 438, Article 227035. <https://doi.org/10.1016/j.jpowsour.2019.227035>
- 11 Li W. Measurement and Correlation of the Solubility of Aminocaproic Acid in Some Pure and Binary Solvents / W. Li, S.C. Du, D.D. Han, S.G. Wu // *Journal of Chemical and Engineering Data*. — 2019. — № 64(12). — P. 5312–5323. <https://doi.org/10.1021/acs.jced.9b00543>
- 12 Kutzhanova K.Z. Nonempirical modeling of protolytic processes in dimeric molecules of amino acids / K.Z. Kutzhanova, A.F. Kurmanova, I.A. Pustolaikina, B.M. Ismagulov // *Bulletin of the University of Karaganda — Chemistry*. — 2018. — № 90. — P. 58–63. <https://doi.org/10.31489/2018ch2/58-63>
- 13 Topol I.A. Protonation of glycine and alanine: proton affinities, intrinsic basicities and proton transfer path / I.A. Topol, S.K. Burt, M. Toscano, N. Russo // *Journal of Molecular Structure — Theochem*. — 1998. — № 430. — P. 41–49.
- 14 Miyamoto H. Allergic contact dermatitis from epsilon-aminocaproic acid / H. Miyamoto, M. Okajima // *Contact Dermatitis*. — 2000. — № 42(1). — P. 50.
- 15 Romero C.M. Effect of Temperature on The Solubility of  $\alpha$ -Amino Acids and  $\alpha,\omega$ -Amino Acids in Water / C.M. Romero, C.D. Oviedo // *Journal of Solution Chemistry*. — 2013. — № 42(6). — P. 1355–1362. doi: 10.1007/s10953-013-0031-9
- 16 Никольский С.Н. Моделирование спектров ЭПР 3,6-ди-*tert*-бутил-2-оксифеноксид в реакции межмолекулярного протонного обмена / С.Н. Никольский // *Вестник ЕНУ — Химия*. — 2007. — № 60(6). — С. 160–167.
- 17 Nikolskiy S.N. Investigation of intermolecular proton exchange of 3,6-di-*tert*-butyl-2-oxyphenoxy with N-phenylanthranilic acid by ESR spectroscopy method / S.N. Nikolskiy, F.Z. Abilkanova, A.S. Golovenko, I.A. Pustolaikina, A.S. Masalimov // *Bulletin of the University of Karaganda-Chemistry*. — 2020. — № 98. — P. 35–41. <https://doi.org/10.31489/2020Ch2/35-41>
- 18 Masalimov A.S. Protolytic Reactions of 3,6-Di-*tert*-Butyl-2-Hydroxyphenoxy with Nitrogen Bases / A.S. Masalimov, A.A. Tur, S.N. Nikolskiy // *Theoretical and Experimental Chemistry*. — 2016. — № 52(1). — P. 57–65. DOI: 10.1007/s11237-016-9451-0

И.Л. Стадник, Ф.Ж. Абилканова, Е.В. Кудрявцева, С.Н. Никольский, А.С. Масалимов

### Толуол ортадағы кейбір алифаттық аминқышқылдарымен протон алмасу реакциясын ЭПР әдісімен зерттеу

Жұмыста толуол сияқты индифферентті еріткіштегі 2-аминогексан қышқылы және 6-аминогексан қышқылы сияқты аминапропн қышқылының кейбір изомерлерімен молекулааралық протон алмасу кинетикасын ЭПР-спектроскопиялық зерттеу нәтижелері келтірілген. Спиндік зонд ретінде 3,6-ди-трет-бутил-2-гидроксифеноксил тұрақты семихинон радикалы пайдаланылды. ЭПР спектрлері РЭ-1306 спектрометрінде тіркелінді. Молекулааралық протон алмасу процесінің алынған динамикалық спектрлері осы мақалада келтірілген. Протон алмасуының ЭПР спектрлерін модельдеу осы реакцияның төрт секірмелі моделіне және модифицирленген Блох теңдеулеріне негізделген бұрын әзірленген бағдарлама арқылы жүзеге асырылды. Модель спектрлерін эксперимент нәтижесінде алынғандармен салыстыру спиндік зондтың аминқышқылдарымен қышқылдық-негіздік әрекеттесуінің кинетикалық параметрлерін ең аз қателікпен анықтауға және реакцияның активтену кедергісін бағалауға мүмкіндік берді. Осылайша алынған мәліметтер талданып, бұрын белгілі ақпаратпен салыстырылды. Зерттелетін алифатты амин қышқылдары спиндік зондпен 3,6-ди-трет-бутил-2-гидроксифеноксилдің алифаттық карбон қышқылдарымен молекулааралық протон алмасу реакциясының жылдамдығымен салыстырылатын жылдамдықпен әрекеттесетіні анықталды. Бірақ, біздің жағдайда, активтену тосқауылының жоғарылауы байқалады, бұл, шамасы, амин қышқылы молекуласындағы амин тобы мен карбоксил тобы арасындағы молекулашілік сутегі байланысы процесіне теріс әсер етумен байланысты.

*Кілт сөздер:* ЭПР спектроскопия, протон алмасу реакциялары, спиндік зонд, семихинонды радикал, Блох теңдеуі, аминогексан қышқылы, 3,6-ди-трет-бутил-2-гидроксифеноксил, ОН-қышқылдары.

И.Л. Стадник, Ф.Ж. Абилканова, Е.В. Кудрявцева, С.Н. Никольский, А.С. Масалимов

### ЭПР-исследование реакции протонного обмена с некоторыми алифатическими аминокислотами в толуольной среде

В работе представлены результаты ЭПР-спектроскопического исследования кинетики межмолекулярного протонного обмена с некоторыми изомерами аминапропновой кислоты, такими как 2-аминогексановая кислота и 6-аминогексановая кислота в среде индифферентного растворителя — толуола. В качестве спинового зонда был использован стабильный семихинонный радикал — 3,6-ди-трет-бутил-2-гидроксифеноксил. Спектры ЭПР регистрировали на спектрометре РЭ-1306. Полученные динамические спектры процесса межмолекулярного протонного обмена приведены в данной статье. Моделирование спектров ЭПР протонного обмена проводилось с помощью ранее разработанной программы, в основе которой лежит четырехпрямая модель данной реакции и модифицированные уравнения Блоха. Сопоставление модельных спектров с экспериментальными позволило нам с минимальной ошибкой определить кинетические параметры процесса кислотно-основного взаимодействия спинового зонда с аминокислотами и оценить активационный барьер реакции. Полученные таким образом данные были проанализированы и сопоставлены с ранее известными сведениями. Было обнаружено, что исследуемые алифатические аминокислоты реагируют со спиновым зондом со скоростями, сопоставимыми со скоростями реакции межмолекулярного протонного обмена 3,6-ди-трет-бутил-2-гидроксифеноксила с алифатическими карбоновыми кислотами. Но, при этом в нашем случае наблюдается рост активационного барьера, который, видимо, связан с негативным влиянием на процесс внутримолекулярной водородной связи между аминогруппой и карбоксильной группой в молекуле аминокислоты.

*Ключевые слова:* ЭПР-спектроскопия, реакции протонного обмена, спиновый зонд, семихинонный радикал, уравнение Блоха, аминогексановая кислота, 3,6-ди-трет-бутил-2-гидроксифеноксил, ОН-кислоты.

#### References

- 1 Wang, C.H., Zhang, R.Q., & Lin, Z.J. (2012). A comparative study on intermolecular hydrogen bond interactions in molecular dimers using different levels of computational methods. *Journal of Theoretical & Computational Chemistry*, 11(6), 1237–1259. <https://doi.org/10.1142/S0219633612500836>
- 2 Bureiko, S.F. (2003, May 25-Jun 01). The kinetic spectroscopy of hydrogen exchange processes in systems with hydrogen bond. Paper presented at the 16th International Conference on Spectroscopy of Molecules and Crystals, Sevastopol, UKRAINE.

- 3 Limbach, H.H., Baumgartner, S., Franke, R., Mannle, F., Scherer, G., & Denisov, G.S. (2021). Double Proton Tautomerism via Intra- or Intermolecular Pathways? The Case of TetramethylReductic Acid Studied by Dynamic NMR: Hydrogen Bond Association, Solvent and Kinetic H/D Isotope Effects. *Molecules*, 26(14), Article 4373. <https://doi.org/10.3390/molecules26144373>
- 4 Grabowski, S. J., Krygowski, T. M., & Stepien, B. (2000). Reaction pathway of proton transfer from the neutral to zwitterionic forms of amino acids. Support for a water molecule-mediated mechanism. *Journal of Physical Organic Chemistry*, 13(11), 740–744. [https://doi.org/10.1002/1099-1395\(200011\)13:11<740::aid-poc304>3.3.co;2-t](https://doi.org/10.1002/1099-1395(200011)13:11<740::aid-poc304>3.3.co;2-t)
- 5 Wu, Z.C., & Fenselau, C. (1992). Proton affinity of arginine measured by the kinetic approach. *Rapid Communications in Mass Spectrometry*, 6(6), 403–405. <https://doi.org/10.1002/rcm.1290060610>
- 6 Bureiko, S.F., & Kucherov, S.Y. (2009). Structure and intermolecular interactions in complexes with hydrogen bond of a series of bifunctional nitrogen-containing compounds. *Journal of Structural Chemistry*, 50(4), 712–721. <https://doi.org/10.1007/s10947-009-0109-9>
- 7 Masalimov, A.S., Kurmanova, A.F., Ospanov, A.U., Tur, A.A., & Nikolskiy, S.N. (2014). EPR-spectroscopy of the fast proton exchange reactions in solutions. *Bulletin of the Karaganda university-Chemistry* (73), 30–35.
- 8 Nikolskiy, S.N., Tur, A.A., Yelchibekova, A.A., Kutzhanova, K.Zh. (2015). Investigation of intermolecular proton exchange 3,6-di-tert-butyl-2-oxyphenoxyl with phenol by ESR spectroscopy method. *Bulletin of the Karaganda university-Chemistry* (77), 47–50.
- 9 Rozman, M. (2012). Proton affinity of several basic non-standard amino acids. *Chemical Physics Letters*, 543, 50–54. <https://doi.org/10.1016/j.cplett.2012.06.048>
- 10 Zhao, G. D., Xu, X. L., Di, Y. B., Wang, H., Cheng, B. W., Shi, L. & Yin, Y. (2019). Amino acid clusters supported by cellulose nanofibers for proton exchange membranes. *Journal of Power Sources*, 438, Article 227035. <https://doi.org/10.1016/j.jpowsour.2019.227035>
- 11 Li, W., Du, S. C., Han, D. D., & Wu, S. G. (2019). Measurement and Correlation of the Solubility of Aminocaproic Acid in Some Pure and Binary Solvents. *Journal of Chemical and Engineering Data*, 64(12), 5312–5323. <https://doi.org/10.1021/acs.jced.9b00543>
- 12 Kutzhanova, K. Z., Kurmanova, A. F., Pustolaikina, I. A., & Ismagulov, B. M. (2018). Nonempirical modeling of protolytic processes in dimeric molecules of amino acids. *Bulletin of the University of Karaganda – Chemistry* (90), 58–63. <https://doi.org/10.31489/2018ch2/58-63>
- 13 Topol, I. A., Burt, S. K., Toscano, M., & Russo, N. (1998). Protonation of glycine and alanine: proton affinities, intrinsic basicities and proton transfer path. *Journal of Molecular Structure – Theochem*, 430, 41–49.
- 14 Miyamoto, H., & Okajima, M. (2000). Allergic contact dermatitis from epsilon-aminocaproic acid. *Contact Dermatitis*, 42(1), 50–50.
- 15 Romero, C. M., & Oviedo, C. D. (2013). Effect of Temperature on The Solubility of  $\alpha$ -Amino Acids and  $\alpha,\omega$ -Amino Acids in Water. *Journal of Solution Chemistry*, 42(6), 1355–1362. doi: 10.1007/s10953-013-0031-9
- 16 Nikolskiy S.N. (2007). Modelirovanie spektrov EPR 3,6-di-tret-butyl-2-oksifenoksila v reakcii mezhmolekulyarnogo protonnogo obmena [Simulation of ESR spectra of 3,6-di-tert-butyl-2-oxyphenoxyl in the reaction of intermolecular proton exchange]. *Vestnik ENU-Himiya — Bulletin of the L.N. Gumilyov Eurasian National University – Chemistry*, 60(6), 160–167 [in Russian].
- 17 Nikolskiy, S.N., Abilkanova, F. Z., Golovenko, A. S., Pustolaikina, I. A., & Masalimov, A. S. (2020). Investigation of intermolecular proton exchange of 3,6-di-tert-butyl-2-oxyphenoxyl with N-phenylanthranilic acid by ESR spectroscopy method. *Bulletin of the University of Karaganda – Chemistry*(98), 35–41. <https://doi.org/10.31489/2020Ch2/35-41>
- 18 Masalimov, A.S., Tur, A.A., & Nikolskiy, S.N. (2016). Protolytic Reactions of 3,6-DI-tert-Butyl-2-Hydroxyphenoxyl with Nitrogen Bases. *Theoretical and Experimental Chemistry*, 52(1), 57–65. <https://doi.org/10.1007/s11237-016-9451-0>

### Information about authors

**Stadnik, Irina Leonidovna** (corresponding author) — Candidate of Chemical Sciences, Assistant Professor, Physical and Analytical Chemistry Department, Karagandy University of the name of academician E.A. Buketov, Universitetskaya street, 28, 100024, Karaganda, Kazakhstan; e-mail: [i.stadnik@mail.ru](mailto:i.stadnik@mail.ru); <https://orcid.org/0000-0002-9497-4177>;

**Abilkanova, Farida Zhumageldyevna** — PhD student, Physical and Analytical Chemistry Department, Karagandy University of the name of academician E.A. Buketov, Universitetskaya street, 28, 100024, Karaganda, Kazakhstan; e-mail: [Farida\\_b88@mail.ru](mailto:Farida_b88@mail.ru); <https://orcid.org/0000-0002-6894-5276>;

**Kudryavtseva, Yelizaveta Vasilyevna** — 1st year master's student, Physical and Analytical Chemistry Department, Karagandy University of the name of academician E.A. Buketov, Universitetskaya street, 28, 100024, Karaganda, Kazakhstan; e-mail: [kudryavcevaev99@gmail.com](mailto:kudryavcevaev99@gmail.com); <https://orcid.org/0000-0002-1672-1318>;

**Nikolskiy, Sergey Nikolayevich** — Doctor of Chemical Sciences, Professor, Physical and Analytical Chemistry Department, Karagandy University of the name of academician E.A. Buketov, Universitetskaya

street, 28, 100024, Karaganda, Kazakhstan; Professor, Physical and Inorganic Chemistry Department, Altay State University, 656049, Barnaul, Russia, e-mail: [sergeynikolsky@mail.ru](mailto:sergeynikolsky@mail.ru); <https://orcid.org/0000-0003-3175-6938>;

**Masalimov, Abay Sabirzhanovich** — Full Professor, Doctor of Chemical Sciences, Physical and Analytical Chemistry Department, Karagandy University of the name of academician E.A. Buketov, Universitetskaya street, 28, 100024, Karaganda, Kazakhstan; e-mail: [masalimov-as@mail.ru](mailto:masalimov-as@mail.ru); <https://orcid.org/0000-0002-6405-7108>